

BASE MANIFOLDS FOR LAGRANGIAN FIBRATIONS ON HYPERKÄHLER MANIFOLDS

DANIEL GREB AND CHRISTIAN LEHN

ABSTRACT. Let $f: X \rightarrow B$ be a fibration from a hyperkähler manifold to a complex space B . Assuming that B is smooth, we show that $B \cong \mathbb{P}^n$. This generalises a theorem of J.-M. Hwang to the Kähler case.

1. INTRODUCTION

One of the most important and startling conjectures in the study of hyperkähler manifolds X says that the base space of any non-trivial fibration $X \rightarrow B$ is the complex projective space $\mathbb{P}^n$, where $n = \dim X/2$. We refer the reader to [GHJ03, 21.4] for a discussion of this conjecture. Any such fibration is automatically Lagrangian with respect to the holomorphic symplectic form by works of Matsushita; see [Mat03] or Section 2 for a summary of his results.

With the additional hypothesis that X be projective and that B be smooth, the conjecture is known to hold by work of Jun-Muk Hwang [Hwa08]. In this short note, we remove Hwang's projectivity assumption on X , and prove the following result:

Theorem 1.1. *Let X be a hyperkähler manifold, and let $f: X \rightarrow B$ be a fibration onto a complex space B . If B is smooth, then $B \cong \mathbb{P}^n$.*

The idea of the proof is simple: after noticing that the base manifold has to be projective, we pull back a very ample line bundle from B to X and use this line bundle to deform the given fibration to a sequence of projective ones. Then, we apply Hwang's theorem to these projective deformations and use global deformation rigidity of $\mathbb{P}^n$ to conclude the desired result for the central fibre.

2. PRELIMINARIES

We start by fixing our notation and by recalling definitions of the basic objects investigated in this note.

Definition 2.1. A compact Kähler manifold is called *hyperkähler* or *irreducible holomorphic symplectic* if it is simply-connected, and if $H^0(X, \Omega_X^2) = \mathbb{C}\sigma$, where σ is everywhere non-degenerate. A *fibration* on X is a surjective holomorphic map $f: X \rightarrow B$ with $f_*\mathcal{O}_X = \mathcal{O}_B$ from X to a complex space B with $0 < \dim B < \dim X$. A *Lagrangian fibration* on X is a fibration $f: X \rightarrow B$ such that every irreducible component of every fibre of f is a Lagrangian subvariety with respect to the holomorphic symplectic form σ .

To make this note more self-contained, we collect some known results concerning fibrations on hyperkähler manifolds, which we will use in the subsequent proof, in the following proposition.

Proposition 2.2. *Let X be a hyperkähler manifold of dimension $2n$, and let $f: X \rightarrow B$ be a fibration onto a complex space B . Then, f is a Lagrangian fibration onto a normal projective variety. In particular, B has dimension n .*

Proof. As explained in [AC13, Thm. 1 and footnote], using results of Varouchas [Var86, Var89] and the fundamental results of Matsushita [Mat99, Mat01, Mat00], one shows without any a priori assumption on the base of the fibration that B is a normal Kähler space. Then, [Mat03, Thm. 2.1 and Thm. 3.1] imply the claim. $\square$

3. PROOF OF THEOREM 1.1

Let X be a hyperkähler manifold of dimension $2n$, and let $f: X \rightarrow B$ be a fibration onto a complex space B .

Since B is projective by Proposition 2.2, there exists a very ample line bundle on B . Let L denote its pullback under f . Furthermore, let $\mathcal{X} \rightarrow (S, 0)$ be the (smooth) Kuranishi space of X ; in particular, $\mathcal{X} \rightarrow (S, 0)$ is a smooth family of hyperkähler manifolds. By [Mat09, Thm. 1.1(1) and 1.1(2)] there exists a smooth hypersurface $(S_L, 0)$ in $(S, 0)$, and a line bundle $\mathcal{L}$ on the pullback $\mathcal{X}_L = S_L \times_S \mathcal{X}$ of $\mathcal{X}$ to S_L such that the restriction of $\mathcal{L}$ to the fibre over the reference point 0 is isomorphic to L . We denote the natural projection $\mathcal{X}_L \rightarrow S_L$ by p , and we note that both $\mathcal{X}_L$ as well as p are smooth. As usual we will take a representative of the germ $(S_L, 0)$ and shrink it if necessary (keeping the base point), usually without mentioning this explicitly.

By [Mat09, Thm. 1.1(3) and Cor. 1.2] the pushforward $p_*\mathcal{L}$ is a vector bundle, the canonical map $p^*p_*\mathcal{L} \rightarrow \mathcal{L}$ is surjective, and thus gives rise to a morphism $F: \mathcal{X}_L \rightarrow \mathbb{P}_{S_L}(p_*\mathcal{L}^\vee)$ over S_L that extends $f: X \rightarrow B$ to the whole family $\mathcal{X}_L$. Let us denote by $\mathcal{B} \subset \mathbb{P}_{S_L}(p_*\mathcal{L}^\vee)$ the scheme-theoretic image of F . We thus have a commutative diagram

$$(3.1) \quad \begin{array}{ccc} \mathcal{X}_L & \xrightarrow{F} & \mathcal{B} \\ & \searrow p & \downarrow \pi \\ & & S_L. \end{array}$$

Lemma 3.1. *We have $b_2(X) \geq 4$. In particular, S_L has positive dimension.*

Proof. Let α be a Kähler class on X . Since L is not ample, its Chern class is not a multiple of α , hence $h^{1,1}(X) \geq 2$, implying the first claim. For the second claim, note that the dimension of the Kuranishi space S of X is $\dim H^1(X, T_X) = h^{1,1}(X) \geq 2$ and S_L is a hyperplane in S . $\square$

So $\mathcal{X}_L \rightarrow S_L$ is a positive-dimensional smooth family of hyperkähler manifolds. Restricting diagram (3.1) to a smooth embedded disk $\Delta \subset S_L$ through the origin, we see that $\mathcal{B}_\Delta = \pi^{-1}(\Delta) \rightarrow \Delta$ is flat with smooth central fibre, hence a smooth morphism. In particular, $\mathcal{B}_\Delta$ is smooth. Moreover, [GHJ03, Prop. 26.6] implies that there exists a dense subset

$T \subset \Delta$ such that $X_t := p^{-1}(t)$ is projective for all $t \in T$. Therefore, by Hwang’s theorem [Hwa08] the fibre $B_t := \pi^{-1}(t)$ is isomorphic to $\mathbb{P}^n$ for all $t \in T$. Hence, we find a sequence of points $\{p_\nu\}_{\nu \in \mathbb{N}}$ in $T \subset \Delta$ such that $\lim_{\nu \rightarrow \infty} (p_\nu) = 0$, and such that $B_t \cong \mathbb{P}^n$. Hence, global deformation rigidity of $\mathbb{P}^n$, see [Siu91, paragraph following the Main Theorem] implies that the central fibre is likewise isomorphic to $\mathbb{P}^n$. This concludes the proof of Theorem 1.1.

Acknowledgements. The authors are grateful to Sönke Rollenske for helpful discussions. The first named author gratefully acknowledges support by the Baden-Württemberg Stiftung through the “Eliteprogramm für Postdoktorandinnen und Postdoktoranden”, as well as by the DFG-Research Training Group GK 1821 “Cohomological methods in geometry”.

REFERENCES

- [AC13] Ekaterina Amerik, Frédéric Campana, *On families of lagrangian tori on hyperkähler manifolds*, arXiv:1303.0613, 2013. – cited on p. 2
- [GHJ03] Mark Gross, Daniel Huybrechts, and Dominic Joyce, *Calabi-Yau manifolds and related geometries*, Universitext, Springer-Verlag, Berlin, 2003, Lectures from the Summer School held in Nordfjordeid, June 2001. MR 1963559 – cited on p. 1, 2
- [Hwa08] Jun-Muk Hwang, *Base manifolds for fibrations of projective irreducible symplectic manifolds*, Invent. Math. **174** (2008), no. 3, 625–644. MR 2453602 – cited on p. 1, 3
- [Mat99] Daisuke Matsushita, *On fibre space structures of a projective irreducible symplectic manifold*, Topology **38** (1999), no. 1, 79–83. MR 1644091 (99f:14054) – cited on p. 2
- [Mat00] ———, *Equidimensionality of Lagrangian fibrations on holomorphic symplectic manifolds*, Math. Res. Lett. **7** (2000), no. 4, 389–391. MR 1783616 (2001f:32041) – cited on p. 2
- [Mat01] ———, *Addendum: “On fibre space structures of a projective irreducible symplectic manifold”*, Topology **40** (2001), no. 2, 431–432. – cited on p. 2
- [Mat03] ———, *Holomorphic symplectic manifolds and Lagrangian fibrations*, Acta Appl. Math. **75** (2003), no. 1-3, 117–123. MR 1975562 (2004c:32040) – cited on p. 1, 2
- [Mat09] Daisuke Matsushita, *On deformations of Lagrangian fibrations*, preprint arXiv:0903.2098, 2009. – cited on p. 2
- [Siu91] Yum Tong Siu, *Global nondeformability of the complex projective space*, Prospects in complex geometry (Katata and Kyoto, 1989), Lecture Notes in Math., vol. 1468, Springer, Berlin, 1991, pp. 254–280. MR 1123546 (93d:32028) – cited on p. 3
- [Var86] Jean Varouchas, *Sur l’image d’une variété kählérienne compacte*, Fonctions de plusieurs variables complexes V (Paris, 1979-1985), Lecture Notes in Math., vol. 1188, Springer, Berlin, 1986, pp. 245–259. MR 0926290 (89d:32064) – cited on p. 2
- [Var89] Jean Varouchas, *Kähler spaces and proper open morphisms*, Math. Ann. **283** (1989), no. 1, 13–52. MR 973802 (89m:32021) – cited on p. 2

DANIEL GREB, ARBEITSGRUPPE ALGEBRA / TOPOLOGIE, FAKULTÄT FÜR MATHEMATIK, RUHR-UNIVERSITÄT BOCHUM, 44780 BOCHUM, GERMANY

E-mail address: daniel.greb@rub.de

URL: <http://www.rub.de/ffm/Lehrstuehle/Greb/index.html>

CHRISTIAN LEHN, INSTITUT DE RECHERCHE MATHÉMATIQUE AVANCÉE, UNIVERSITÉ DE STRASBOURG, 7 RUE RENÉ DESCARTES, 67084 STRASBOURG CEDEX, FRANCE

E-mail address: lehn@math.unistra.fr